

OPIC
OFFICE DE LA PROPRIÉTÉ
INTELLECTUELLE DU CANADA



CIPO
CANADIAN INTELLECTUAL
PROPERTY OFFICE

Ottawa Hull K1A 0C9

(11) (C)	2,054,898
(22)	1991/11/04
(43)	1992/05/08
(45)	1996/12/03

(51) Int.Cl. ⁶ C11D 1/94; A61K 7/075; A61K 7/50

(19) (CA) **CANADIAN PATENT** (12)

(54) Detergent Composition

(72) Edwards, Robert James , Germany (Federal Republic of)
Lee, Robert Stanley , U.K.
Salmon, Tom Matthew Forrest , U.K.

(73) Unilever PLC , U.K.

(30) (GB) U.K. 9024162.1 1990/11/07

(57) 27 Claims

NO DRAWING



Industrie
Canada

Industry
Canada

OPIC - CIPO 101

Canada

03 DEC. 1996

ABSTRACTDETERGENT COMPOSITION

A foaming detergent composition for personal washing comprises:

A - a detergent active, preferably a mixture of

A1 - non-soap anionic detergent, and

A2 - amphoteric or polar nonionic detergent; and

B - a skin emollient material, preferably selected from

B1 - skin moisturizers

B2 - occlusive agents

B3 - humectants

and mixtures thereof.

DETERGENT COMPOSITION

This invention is concerned with liquid detergent compositions suitable for personal washing. Examples of
5 such compositions are shower gels and bath foams. A requirement for such compositions is that they should generate a lather when used.

Emollient materials are typically used in a skin cream to help restore and maintain the water content of the
10 skin. At least some emollient materials are inherently antifoaming in nature and therefore appear unsuitable for incorporation in a product which is required to generate a lather in use.

We have, however, now found that it is possible
15 to incorporate such materials into a liquid product.

Broadly, therefore, this invention provides an aqueous liquid detergent composition containing foaming detergent active and skin emollient material.

A - Detergent Active Materials

20 The detergent active materials employed in the present invention are selected from the classes consisting of anionic, amphoteric, and nonionic detergents. It is



preferred that the detergent active is a mixture of:

A1 - non-soap anionic detergent, and

A2 - amphoteric or polar nonionic detergent.

5 Preferred anionic detergents (A1) are alkyl ether
sulphates, acyl isethionates, alkyl glycerol ether
sulphonates, acyl glutamates, acyl peptides, sarcosinates,
ester carboxylic acids, α -olefin
sulphonates, sulphosuccinates, alkyl benzene sulphonates,
10 amides of N-methyl taurine, α -sulpho fatty acids and alkyl
sulphates. Such detergents generally include an alkyl or
acyl group of 8 to 20 carbon atoms, together with an
anionic group.

It is particularly preferred that the second
15 detergent (A2) is an amphoteric surfactant which may be a
betaine, a sulphobetaine or an amido betaine. All of these
are zwitterionic. Less preferred are acyl ethanolamides,
acyl diethanolamides and amine oxides, all of which are
polar nonionic surfactants.

20 B - Skin Emollient Agents

These materials may be chosen from three
categories which are

B1 - Skin moisturisers

B2 - Occlusive agents

B3 - Humectants

Skin moisturisers (B1) are compounds which
5 enhance moisture retention by the skin and may also attract
water into the skin. Typical moisturising materials
include urea, lactic acid, pyrrolidone carboxylic acid,
amino acids and salts of the acids mentioned.

The occlusive agents (B2) are substances which
10 form on the skin thin films of limited permeability,
serving to hold water within the skin and prevent
dehydration. The range of occlusive agents is
considerable. They are generally hydrophobic oils and
waxes. Examples of classes of such agents and individual
15 examples of such agents are:

1. Hydrocarbon oils and waxes. Examples thereof
are mineral oil, petrolatum, paraffin, ceresin, ozokerite,
microcrystalline wax.

2. Silicone oils, such as dimethyl
20 polysiloxanes, methylphenyl polysiloxanes, silicone glycol
copolymers.

3. Triglyceride esters, for example vegetable
and animal fats and oils.

4. Glyceride esters and esters such as acetylated monoglycerides, and ethoxylated monoglycerides.

5. Alkyl and alkenyl esters of fatty acids having 10 to 20 carbon atoms. Examples include hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl myristate, isopropyl palmitate, decyl oleate, isodecyl oleate, hexadecyl stearate, decyl stearate, isopropyl isostearate, diisopropyl adipate, diisohexyl adipate, dihexyldecyl adipate, diisopropyl sebacate, lauryl lactate, myristyl lactate, cetyl lactate, oleyl myristate, oleyl stearate and oleyl oleate.

6. Fatty alcohols having 10 to 20 carbon atoms. Lauryl, myristyl, cetyl, hexadecyl, stearyl, isostearyl, hydroxystearyl, oleyl, ricinoleyl, behenyl, erucyl, and 2-octyl dodecanyl alcohols are examples of satisfactory fatty alcohols.

7. Lanolin and derivatives. Lanolin, lanolin oil, lanolin wax, lanolin alcohols, lanolin fatty acids, isopropyl lanolate, ethoxylated lanolin, ethoxylated lanolin alcohols, ethoxylated cholesterol, propoxylated lanolin alcohols, acetylated lanolin, acetylated lanolin alcohols and lanolin alcohols linoleate are illustrative emollients derived from lanolin.

8. Natural waxes, esters thereof and ethoxylated natural waxes, beeswax, spermaceti, myristyl myristate, stearyl stearate, polyoxyethylene sorbitol beeswax, carnauba wax and candelilla wax.

5 Especially preferred are C_2 - C_{18} alkyl esters of C_{12} - C_{18} fatty acids, such as isopropyl myristate, and isopropyl palmitate.

Humectants (B3) are especially C_2 - C_6 polyols notably glycerol, sorbitol, propylene glycol and 1,3-
10 butylene glycol. A further example of humectant is polyethylene glycols having molecular weights of from about 100 to about 1500. Humectants do not themselves form occlusive films but may cooperate with other materials to form a film having occlusive properties. It is therefore
15 desirable that humectants are not the sole category of skin emollient agent present.

The overall amount of emollient agents included in a composition of this invention is preferably in a range from 0.1 to 10% of the composition more preferably 0.2 to
20 5% by weight of the composition. Somewhat more specifically, it is preferred that the skin emollient agents include occlusive agent (B2) in an amount from 0.1 to 2% by weight of the composition, better 0.2 to 1%. If skin moisturiser (B1) is present the amount is preferably
25 0.02 to 5%, better 0.05 to 2% of the composition. If

humectant (B3) is present the amount will generally not exceed 15% by weight of the composition and it will generally be convenient for humectant to be present in a quantity from 0.1 to 15% better 0.5 to 5 or 8% by weight of the composition.

Particularly preferred weight ratios of the three categories of skin emollient agent are a weight ratio of

$$B1:B2:B3 = 0-15:2-10:5-50$$

Preferred amounts of the detergent active lie in a range from 3-40% by weight of the overall composition, preferably 5 to 25% by weight. Preferably anionic detergent active (A1) is present in an amount from 5 to 20% by weight of the composition while a second detergent active (A2) is present in an amount from 0.5 to 15%, preferably 1 to 10% by weight of the composition. The two categories of detergent active A1:A2 are preferably in a weight ratio from 10:1 to 1:10, better 9:1 to 1:2.

The weight ratio of detergent active to skin emollient agents preferably lies in a range from 20:1 to 5:1. Alternatively or additionally the weight ratio of detergent active to occlusive agent lies in a range from 10:1 to 100:1.

Thus, a preferred form of this invention comprises the materials A1, A2 and B1-B3 as explained above

in amounts by weight which are:

	A1	:	5-20%
	A2	:	0.5-15%
	B1	:	0.02-5%
5	B2	:	0.1-2%
	B3	:	0-15% preferably 0.5 to 8%

Other materials may be included in compositions according to this invention. In particular, electrolyte or thickening agents or both may be incorporated to control the viscosity of the composition. We prefer that compositions of this invention should have a viscosity in the range from 10,000 to 1000 mPs at 10 sec^{-1} shear rate, more preferably 8000 to 2000 mPs.

Thickening agents may be synthetic cross-linked polymers such as cross-linked polyacrylates. Another category of thickening agents are polysaccharides such as vegetable mucins, alginates, cellulose derivatives and xanthan gums. The amount of any thickening agent is likely to range from 0.01 to 5% by weight of the composition and likely narrower limits are 0.05 to 2% by weight of the composition.

Further materials which are likely to be included are pearlescers or opacifiers, perfumes, colorants and antioxidants. Preservatives may well be included.

Free fatty acids typically of 8 to 22 carbon atoms may be present or may be avoided. Soaps are possible but are preferably avoided. If soap or fatty acid is present the amount of each is unlikely to exceed 5% by weight of the composition and more particularly may be in an amount from 0.5 to 2.5% by weight of the composition.

One feature which is possible within this invention is that the skin emollient agents (B) may be capable of incorporation at the same relative proportions into a stable emulsion. The concentration of these agents in such an emulsion would be the same or more probably greater than their concentration in the detergent composition of the invention.

This feature creates the possibility of incorporating the formulation of one product form, namely a skin moisturising liquid or cream, into another product form, namely the liquid detergent composition of the invention.

When the invention takes such a form it can be defined as an aqueous liquid detergent composition containing

foaming detergent active and
the non-aqueous constituents of a moisturising emulsion which comprises skin emollient agents.

2054898

Such a moisturising emulsion might include other materials which are also incorporated into the detergent composition, for example thickeners, opacifiers or pearlescers.

5 Forms of the invention which are of special interest are those in which the detergent active comprises C_8 to C_{12} fatty acyl isethionate. Fatty acyl isethionate is known as mild to the skin but is poorly soluble. We have observed that combinations of fatty acyl isethionate
10 with an amphoteric surfactant or a polar nonionic surfactant can give solutions which are clear and isotropic (if in the absence of opacifier, pearlescer or other non-detergent insolubles).

Consequently forms of this invention can be
15 defined as detergent compositions containing

- A) 3 to 40% of detergent composed of
 - A1) at least 2% of C_8 - C_{12} acyl isethionate
 - A2) at least 1% of amphoteric or polar nonionic detergent
- B) 0.1 to 10% of skin emollient agents

20 wherein the ratio of acyl isethionate to detergent A2 is from 10:1 to 1:4, better 5:1 to 1:2, and the detergent actives present in the composition form a clear solution in distilled water at the same concentrations.

The detergent actives present may include

additional anionic detergent (A1) and the amount may be such as to keep the ratio of total anionic (A1) to second detergent (A2) within a range of 10:1 to 1:2.

Method of Preparation

5 The compositions of this invention can be prepared by simply mixing their constituents with water. Premixes of some components may be made and combined, if convenient. It will generally be desirable to arrange the order of addition so that thickening occurs after most of
10 the mixing has taken place.

Example 1

Shower gels were prepared with formulations as set out in Table 1 below.

15 In each case the anionic detergent (A1) was sodium lauryl ether sulphate (SLES) with an average of three ethylene oxide groups per molecule. The second detergent (A2) was coconut amido propyl betaine (CAPB) opacifier was a polystyrene latex. The alternative to this was a pearlescent agent which was ethylene glycol
20 distearate (EGDS).

"NMF" denotes a mixture of materials sold as equivalent to a natural moisturising factor in skin. Such

a mixture is available commercially as HYDROVITON (trade mark) from Dragoco, LACTIL (trade mark) from Goldsmith or PRODEW (trade mark) from Ajinomoto. The NMF used contained 40% of free amino acids, 12% pyrrolidone carboxylic acid, 7% urea and 12% lactate, the balance being made up by other constituents. Thus this material contained at least 70% of skin moisturising agents.

TABLE 1

<u>Formulation:</u>		A	B	C	D
10	Ingredients	% by weight			
	SLES	12	12	12	12
	CAPB	3	3	3	3
	Opacifier	1	1	-	-
	Pearlescer	-	-	4	4
15	Isopropyl palmitate	-	0.5	-	0.5
	Glycerol	-	1.25	-	1.25
	NMF	-	0.25	-	0.25
	Preservative	0.07	0.07	0.07	0.07
	Sodium chloride	4.0	4.0	4.0	4.0
20	Water	- - balance to 100% --			

The four formulations were assessed by a panel of ten persons. Each panelist wore surgical gloves. The gloves were first washed with a standard detergent composition in running water at 30°C. One gram of the test composition was then dispensed from a syringe into the

palm of each panelist. The panelists were asked to wash their hands as they would normally, using water of 26° French hardness (FH) at 30°C. The panelists were asked to score the lather volume and perceived creaminess of the lather. The actual amount of lather produced was assessed by collecting the lather in an inverted funnel which is then used to transfer the lather to a graduated cylinder.

The ability of the compositions to generate foam was also determined by the following test. 5 grams of the formulation was put into a graduated separating funnel. 50 grams of water is then added. This water was of 26° FH at 25°C. The funnel is shaken for 10 seconds and the foam level measured and noted as "initial foam". The solution below the foam is drained off, a further 50 grams of water is added, the funnel is shaken again and foam volume noted. The draining off of solution, addition of water and shaking is repeated until all foam has disappeared. The number of additions of 50 gram aliquots of water is referred to as "number of rinses" and is an indication of the amount of water required to rinse the foam. Each time an aliquot of water was added and the separating funnel shaken the amount of foam present was noted. The total of these amounts of foam is referred to as the cumulative foam volume.

Table 2 sets out total of normalised panel scores, cumulative foam volume and number of rinses (i.e.

number of aliquots added until no foam remains) for each of the four formulations.

The results show that the inclusion of emollient agents in formulations B and D had little or no adverse effect on foaming. In the case of compositions A and B, the panel test and funnel test give opposite results but it can be seen that the effect of including the emollient agents is much less than the change from opacifier to pearlescer.

TABLE 2

Formulation:	A	B	C	D
Total Normalised Panel Scores:				
Lather Volume	142.7	102.3	98.1	102.1
5 Lather Creaminess	111.7	105.4	103.9	99.2
<u>Funnel Test</u>				
Cumulative Foam Volume	6550	6900	2025	2550
No Rinses to Zero Foam	15	13	11	11

Example 2

Creams were prepared using emollient agents and pearlescer or opacifier as used in Example 1. Crosslinked polyacrylate was included. The formulations were as

5 follows:

<u>Formulation:</u>		<u>E</u>	<u>F</u>
Ingredient		wt%	wt%
10	Glycerol	5	5
	Isopropyl palmitate	2	2
	NMF	1	1
	Pearlescer	16	-
	Opacifier	-	4
	Perfume	1	1
	Polyacrylate	1	1
15	Water	-- balance to 100% --	

The method of producing the creams was to mix all of the materials with water, incorporating the polyacrylate in the acid form supplied by the manufacturers, and shaken to form an emulsion. The resulting compositions were

20 viscous stable creams.

Example 3

Liquid compositions were prepared generally as in Example 1 but incorporating a sulphosuccinate in place of

betaine. One composition incorporated 25% of glycerol.

The formulations were:

	<u>Formulation:</u>	G	H
	Ingredient	% by weight	
5	SLES	10	10
	Sulphosuccinate	2	2
	Tegobetaine	3	3
	Glycerol	-	25
	Perfume	1.5	1.5
10	Sodium chloride	1.5	1.5

The compositions were assessed by panel testing as in Example 1 and the results obtained were as follows:

		G	H
15		(with glycerol)	
	Panel Scores:		
	Lather Volume	158	165
	Lather Creaminess	125	148

Example 4

20 Example 3 was repeated, but one formulation contained 10% of sodium lactate. The results were as follows:

17

Total Panel Scores	without lactate	with lactate
Lather Volume	227	188
Lather Creaminess	162	174

These Examples 3 and 4 show that up to 25%
5 glycerol and up to 10% lactate can be included without
serious adverse effect.

Example 5

A shower gel was prepared generally as Example
1D, but containing a different detergent active system,
10 which includes sodium cocoyl isethionate.

The foaming ability of this composition was
compared with two compositions similar to Examples 1C and
1D. The three formulations were as set out in Table 4
below. SLES, CAPB, NMF and pearlescer are the same
15 materials as used in Example 1.

The test procedure was as follows. 2 gram of
test composition are dissolved in 20mls of distilled water.
20mls of this solution are put into a 100ml stoppered
measuring cylinder. The cylinder is shaken 10 times,
20 swapped to the other hand and shaken 10 times again. The
height of the lather is measured immediately. The
operation is performed twice for each product, and the
results are an average of the two measurements.

The data obtained by this method are included in
Table 4 below.

TABLE 4				
<u>Formulation:</u>		<u>I</u>	<u>J</u>	<u>K</u>
5	Ingredients	% by weight		
	Sodium cocoyl isethionate	5	-	-
	SLES	2	13	13
	CAPB	8	2	2
	Pearlescer	4	4	4
10	Isopropyl palmitate	0.5	-	0.5
	Glycerol	1.25	-	1.25
	NMF	0.25	-	0.25
	Preservative	0.07	0.07	0.07
	Sodium chloride	3.5	4.0	4.0
15	Water	-- balance to 100% --		
	Foam height (cm)	14.7	12.5	15.4

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A detergent composition in the form of an aqueous liquid of viscosity in the range from 10,000 to 1,000 mPas at 10 sec⁻¹ shear rate, comprising:

A - foaming detergent active which is a mixture of:

A1 - at least 5% by weight of the composition of a non-soap anionic detergent which is selected from alkyl ether sulphates, acyl isethionates, alkyl glycerol ether sulphonates, acyl glutamates, acyl peptides, sarcosinates, ester carboxylic acids, α -olefin sulphonates, sulphosuccinates, alkyl benzene sulphonates, amides of N-methyl taurine, α -sulpho fatty acids and alkyl sulphates; with the proviso that said anionic detergent includes at least 2% by weight of the composition of C₈-C₂₂ acyl isethionate;

A2 - from 1 to 15% by weight of the composition of amphoteric detergent selected from betaines, sulphobetaines and amidobetaines; and

B - at least 0.1% by weight of a skin emollient material.

2. A detergent composition according to claim 1, wherein the skin emollient material (B) is selected from:

B1 - skin moisturisers

B2 - occlusive agents

B3 - humectants

and mixtures thereof.

3. A detergent composition according to claim 2, wherein the skin moisturiser, if present is selected from urea, lactic acid, pyrrolidone carboxylic acid, amino acids, and salts of these acids.

4. A detergent composition according to claim 2, wherein the occlusive agent, if present, is selected from: hydrocarbon oils and waxes; silicone oils; triglyceride esters; glyceride esters, acetylated monoglycerides, alkoxyated monoglycerides; alkyl and alkenyl esters of fatty acids having 10 to 20 carbon atoms; fatty alcohols having 10 to 20 carbon atoms; lanolin and derivatives thereof; natural waxes, esters thereof and alkoxyated natural waxes.

5. A detergent composition according to claim 2, wherein the humectant, if present, is selected from C₂-C₆ polyols; polyethylene glycols having molecular weights of from about 100 to about 1500.

6. A detergent composition according to claim 1 or claim 2, wherein the skin emollient material (B) is present in an amount of from 0.1 to 10% by weight of the composition.

7. A detergent composition according to claim 2, wherein:
the skin moisturiser (B1), if any, is present in an amount of from 0.02 to 5% by weight, of the composition; the

occlusive agent (B2), if any, is present in an amount of from 0.1 to 2% by weight of the composition; and the humectant (B3), if any, is present in an amount of up to 15% by weight of the composition.

5

8. A detergent composition according to claim 7, wherein the weight ratios of the three categories of skin emollient material are B1:B2:B3 = 0-15:2-10:5-50.

9. A detergent composition according to claim 1 or claim 2, wherein the detergent active (A) is present in an amount not exceeding 40% by weight of the composition.

10. A detergent composition according to claim 1, wherein the non-soap anionic detergent (A1) is present in an amount of from 5 to 20% by weight of the composition, and the amphoteric detergent (A2) is present in an amount of from 1 to 15% by weight of the composition.

11. A detergent composition according to claim 10, wherein the weight ratio of the two categories of detergent active is A1:A2 = 10:1 to 1:10.

12. A detergent composition according to claim 10, wherein the weight ratio of the two categories of detergent active is A1:A2 = 10:1 to 1:2.

13. A detergent composition according to claim 1 or claim

2, wherein the weight ratio of foaming detergent active (A) to skin emollient material (B) is A:B = 20:1 to 5:1.

14. A detergent composition according to claim 2, wherein the weight ratio of foaming detergent active (A) to occlusive agent (B2) is A:B2 = 10:1 to 100:1.

15. A detergent composition according to claim 2, wherein the skin emollient material comprises a humectant (B3).

16. A detergent composition according to claim 2, wherein the skin emollient material comprises a humectant (B3) together with a skin moisturiser (B1).

17. A detergent composition according to claim 2, wherein the skin emollient material comprises a humectant (B3) together with an occlusive agent (B2).

18. A detergent composition according to claim 2, wherein the skin emollient material comprises:

B1 - 0.02-5% of a skin moisturiser,

B2 - 0.1-2% of an occlusive agent; and

B3 - 0-15% of a humectant.

19. A detergent composition according to claim 1 or claim 2, wherein the weight ratio of acyl isethionate (A1) to detergent (A2) is from 10:1 to 1:4 and the detergent actives present in the composition form a clear solution

in distilled water at the same concentrations.

20. A detergent composition in the form of an aqueous liquid of viscosity in the range from 10,000 to 1,000 mPas at 10 sec⁻¹ shear rate, comprising:

A - foaming detergent active which is a mixture of:

A1 - at least 5% by weight of the composition of a non-soap anionic detergent which is selected from alkyl ether sulphates, acyl isethionates, alkyl glycerol ether sulphonates, acyl glutamates, acyl peptides, sarcosinates, ester carboxylic acids, α -olefin sulphonates, sulphosuccinates, alkyl benzene sulphonates, amides of N-methyl taurine and α -sulpho fatty acids;

A2 - at least 1% by weight of the composition of amphoteric detergent selected from betaines, sulphobetaines and amidobetaines; and

B - a skin emollient material which is a mixture of:

B1 - 0.02-5% of skin moisturiser,

B2 - at least 0.1% of occlusive agent; and

B3 0-15% of humectant.

21. A detergent composition according to claim 20, wherein the skin moisturiser is selected from urea, lactic acid, pyrrolidone carboxylic acid, amino acids, and salts of these acids.

22. A detergent composition according to claim 20, wherein the occlusive agent is selected from:

hydrocarbon oils and waxes; silicone oils; triglyceride esters; glyceride esters, acetylated monoglycerides, alkoxyated monoglycerides; alkyl and alkenyl esters of fatty acids having 10 to 20 carbon atoms; fatty alcohols having 10 to 20 carbon atoms; lanolin and derivatives thereof; natural waxes, esters thereof and alkoxyated natural waxes.

23. A detergent composition according to claim 20, wherein the humectant, if present, is selected from C_2-C_6 polyols; polyethylene glycols having molecular weights of from about 100 to about 1500.

24. A detergent composition according to claim 20 wherein the humectant (B3) is present in an amount of 0.1 to 15% by weight of the composition.

25. A detergent composition according to claim 20, wherein the weight ratio of the two categories of detergent active is $A1:A2 = 10:1$ to $1:2$.

26. A detergent composition according to claim 20 or claim 25, wherein the weight ratio of foaming detergent active (A) to skin emollient material (B) is $A:B = 20:1$ to $5:1$.

27. A detergent composition according to claim 1, claim 2 or claim 20, further comprising one or more additional

- 25-

2054898

materials selected from thickening agents, pearlescers or
opacifiers, perfumes, colourants, antioxidants and
preservatives.



SUBSTITUTE
REMPLACEMENT

SECTION is not Present
Cette Section est Absente
